Sickness & DiarrhoeaUntil vomiting and diarrhoea have stoConjunctivitis (Eye infection)Until discharge has stopped, or treatment	
	nent commenced.
Scabies Until successfully treated	
Impetigo Until lesions are crusted or healed, or	48 hours after treatment has commenced
Ringworm Until treatment has started	
Ear infection Until treatment has commenced for 24	4 hours and child is well
Throat/Chest Infection Child should be away for 2.3 days unt	til treated & visibly well
Chicken Pox Child should be away until blisters have	ve dried up
Measles 7 - 10 days from onset of rash	
Rubella (German Measles) 4 days from onset of rash	
Mumps Until swelling has gone	
Whooping Cough 21 days from onset of cough	
Scarlet Fever Child can return back when been on A	Anti-biotics for 3 - 4 days & no longer have a rash
Hand, Foot & Mouth Child should be kept away until they a	are feeling better
Head lice Until treated	
Anti biotics Off nursery for 24 hours from initial do	se of antibiotics
High TemperatureChild is require to stay at home for 24	hours if they have a high temperature.

HOSPITAL TREATMENT	ABSENCE FROM DAYCARE
Tonsils & Adenoids	One week from operation
Grommets Only	Two days from operation
Asthma Attack	Twenty Four Hours from discharge
Other Hospital treatment	As advised by Doctor

The guide lines above are set by the infection control team at the P.E.H Please respect the advice given as Little Angels endeavour at all times to reduce the spread of infection .